




MALNUTRITION QUALITY
IMPROVEMENT INITIATIVE


Overview of MQii's Electronic Clinical Quality Measures (eCQMs) & Suggested Quality Indicators

The Malnutrition Quality Improvement Initiative (MQii) is a project of the Academy of Nutrition and Dietetics, Avalere Health, and other stakeholders who provided guidance and expertise through a collaborative partnership. Support provided by Abbott.

Malnutrition Electronic Clinical Quality Measures (eCQMs) Align with the Malnutrition Care Workflow



These four developed quality measures help providers understand how they are performing against quality improvement goals set forth in the MQii Toolkit

 = Measure developed to address this step in the malnutrition care workflow

Measure Specification Outline

THE TABLE BELOW OUTLINES THE KEY COMPONENTS NECESSARY TO IMPLEMENT EACH OF THE QUALITY MEASURES IN THE HOSPITAL SETTING AND THE INFORMATION THAT APPEARS ON THE FOLLOWING SLIDES

NQF #: Measure Title

Initial Population	Description of all patients to be evaluated by the measure who share a common set of specified characteristics
Numerator	Describes the process, condition, event, or outcome that satisfies the measure focus or intent.
Denominator	Describes the population evaluated by the measure. The target population defined by the denominator can be the same as the initial population or can be a subset of the initial population to further constrain the population for the purpose of the measure.
Denominator Exclusions	Patients who should be removed from the measure population and denominator before determining if numerator criteria are met.
Data Source	The primary source document(s) used for data collection (for example, billing or administrative data, encounter form, enrollment forms, medical record).
Data Elements	Specific data elements required to fulfill measure specifications and assign patient episodes into denominator and numerator populations. For example, data elements derived from the EHR will incorporate value sets of coding terminology (ICD-9, SNOMED, etc.) that represent the clinical data or process being documented, these must be programmed into the EHR in advance.

Malnutrition Measure #1 Captures the Completion of a Malnutrition Screening within 24 Hours

NQF #3087: Completion of a Malnutrition Screening Within 24 hours of Admission

Initial Population	All patients age 18 years and older at time of admission who are admitted to an inpatient hospital
Numerator	<p>Patients in the denominator who have a completed malnutrition screening documented in the medical record within 24 hours of admission to the hospital. For the purposes of this measure, it is recommended that a malnutrition screening be performed using a validated screening tool which may include but is not limited to one of the following validated tools:</p> <p>Malnutrition Screening Tool (MST) (Wu, 2012), Nutrition Risk Classification (NRC) (Kovacevich, 1997), Nutritional Risk Index (NRI) (Honda, 2016), Nutritional Risk Screening 2002 (NRS-2002) (Bauer, 2005), Short Nutrition Assessment Questionnaire (SNAQ) (Pilgrim, 2016).</p>
Denominator	All patients age 18 years and older at time of admission who are admitted to an inpatient hospital
Denominator Exclusions	Patients with a length of stay of less than 24 hours
Data Source	Electronic Health Record
Data Elements	Inpatient Admission (Time Stamp); Length of Stay (Calculated); Completion of Malnutrition Screening; Interval, in hours, between Malnutrition Screening and Inpatient Admission (Calculated)

Measure Rationale & Clinical Guidance

RATIONALE FOR THIS MEASURE AND THE CLINICAL GUIDANCE SUPPORTING IT ARE OUTLINED BELOW:

Measure Rationale:

- Patients who are malnourished while in the hospital have been associated with important adverse patient safety outcomes such as increased risk of complications, readmissions, and length of stay. Patients who experience these increased risks are also associated with a significant increase in costs. Malnutrition is also associated with many adverse outcomes including depression of the immune system, impaired wound healing, muscle wasting, and increased mortality.
- Screening for the risk of malnutrition in care settings is important to enable early and effective interventions for patients who are malnourished or at-risk of malnutrition. These screenings are the first step in providing optimal, evidence-based malnutrition care for patients.
- Although a review of nationally-representative data on cost and utilization indicated that in 2010, 3.2% of patients had a diagnosis of malnutrition (Corkins, 2014), this may be a severely underreported figure identified in other research studies which have estimated that 4-19 million cases are left undiagnosed and untreated.

Clinical Guidance Support:

- The American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.) recommends the following: Screening for nutrition risk is suggested for hospitalized patients (Evidence Grade E)

Malnutrition Measure #2 Captures the Completion of a Nutrition Assessment for At-Risk Patients

NQF #3088: Completion of a Nutrition Assessment for Patients Identified as At-Risk for Malnutrition within 24 hours of a Malnutrition Screening

Initial Population Patients age 65 years and older at time of admission who are admitted to an inpatient hospital

Numerator Patients in the denominator who have a nutrition assessment documented in the medical record within 24 hours of the most recent malnutrition screening

Denominator Patients who were identified as at-risk for malnutrition upon completing a malnutrition screening

Denominator Exclusions Patients with a length of stay of less than 24 hours

Data Source Electronic Health Record

Data Elements Inpatient Admission (Time Stamp); Length of Stay (Calculated); Malnutrition Screening Result "At-Risk"; Completion of a Nutrition Assessment; Interval, in hours, between Malnutrition Screening result and Nutrition Assessment result (Calculated)

Measure Rationale & Clinical Guidance

RATIONALE FOR THIS MEASURE AND THE CLINICAL GUIDANCE SUPPORTING IT ARE OUTLINED BELOW:

Measure Rationale:

- Referral rates for dietetic assessment and treatment of malnourished patients have proven to be suboptimal, thereby increasing the likelihood of developing such aforementioned complications. Although a review of nationally-representative data on cost and utilization indicated that in 2010, 3.2% of patients had a diagnosis of malnutrition, this may be a severely underreported figure identified in other research studies which have estimated that 4-19 million cases are left undiagnosed and untreated.
- Patel et al. (2014) conducted a national survey of hospital-based professionals in the United States focused on nutrition screening and assessment practices and associated gaps in knowledge of nutrition care. Out of 1,777 unique respondents, only 23.1% reported using a validated assessment tool to help identify clinical characteristics for a malnutrition diagnosis.
- Nutrition assessments conducted for at-risk patients identified by malnutrition screening using a validated screening tool was associated with key patient outcomes including less weight loss, reduced length of stay, improved muscle function, better nutritional intake, and fewer readmissions.

Clinical Guidance Support:

- The American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.) recommends the following: Screening for nutrition risk is suggested for hospitalized patients (Evidence Grade E); Nutrition assessment is suggested for all patients who are identified to be at nutrition risk by nutrition screening (Evidence Grade E)

Malnutrition Measure #3 Ensures Appropriate Malnutrition Diagnosis for Those Malnutrition Findings

NQF #3090: Appropriate Documentation of a Malnutrition Diagnosis

Initial Population

Patients age 65 years and older admitted to inpatient care who have a completed nutrition assessment with findings of malnutrition documented in their medical record

Numerator

Patients with a documented diagnosis of malnutrition

Denominator

Patients age 65 years and older admitted to inpatient care who have a completed nutrition assessment with findings of malnutrition documented in their medical record

Denominator Exclusions

Patients with a length of stay of less than 24 hours; Patients who left against medical advice (AMA); Patients discharged to hospice care

Data Source

Electronic Health Record and Chart Abstracted Clinical Data

Data Elements

Inpatient Admission (Time Stamp); Length of Stay (Calculated); Completed Nutrition Assessment; Nutrition Assessment Findings (Chart Abstracted); Malnutrition Diagnosis

Measure Rationale & Clinical Guidance

RATIONALE FOR THIS MEASURE AND THE CLINICAL GUIDANCE SUPPORTING IT ARE OUTLINED BELOW:

Measure Rationale:

- Data analyzed from the Healthcare Cost and Utilization Project (HCUP), a nationally-representative data set describing U.S. hospital discharges, indicated that approximately 3.2% of hospital discharges included malnutrition as a diagnosis in 2010. However, as this same research article reported, past studies have used validated screening tools to indicate a substantially higher prevalence of malnutrition that has gone undiagnosed in the hospital ranging from 33% to 78%.
- Patients who are malnourished while in the hospital have been associated with important negative outcomes such as increased risk of complications, readmissions, and length of stay. Malnutrition is also associated with many adverse outcomes including depression of the immune system, impaired wound healing, muscle wasting, and increased mortality.
- Nutritional status and progress are often not adequately documented in the medical record. It can be difficult to tell when (or if) patients are consuming food and supplements. In addition, nutritional procedures and EHR-triggered care are often lacking in the hospital. The current evidence supports the early and rapid identification of malnutrition status in order to allow for timely treatment of malnutrition in the hospital. Part of the recommended process for implementing nutrition care is appropriate recognition of the nutrition status, diagnosis, and documentation of that status and diagnosis to address their condition with an appropriate plan of care and communicate patient needs to other care providers.

Clinical Guidance Support:

- A.S.P.E.N. recommends the completion of a nutrition assessment as a comprehensive approach to diagnosing nutrition problems such as malnutrition in order to form the basis for an appropriate nutrition intervention. (Mueller, 2011)

Malnutrition Measure #4 Captures the Documentation of a Nutrition Care Plan for Malnourished Patients

NQF #3089: Nutrition Care Plan for Patients Identified as Malnourished after a Completed Nutrition Assessment

Initial Population	Patients age 65 years and older admitted to inpatient care who have a completed nutrition assessment documented in their medical record
Numerator	Patients with a nutrition care plan documented in the patient's medical record. Care plan components include, but are not limited to: Completed assessment results; data and time stamp; treatment goals; prioritization based on treatment severity; prescribed treatment/intervention; identification of members of the Care Team; and timeline for patient follow-up
Denominator	Patients from the initial population with completed nutrition assessment documented in their medical record with findings of malnutrition
Denominator Exclusions	Patients with a length of stay of less than 24 hours
Data Source	Electronic Health Record and Chart Abstracted Clinical Data
Data Elements	Inpatient Admission (Time Stamp); Length of Stay (Calculated); Completed Nutrition Assessment; Nutrition Assessment Findings (Chart Abstracted); Documented Nutrition Care Plan (Chart Abstracted)

Measure Rationale & Clinical Guidance

RATIONALE FOR THIS MEASURE AND THE CLINICAL GUIDANCE SUPPORTING IT ARE OUTLINED BELOW:

Measure Rationale:


- Referral rates for nutrition assessment and treatment of malnourished patients have proven to be suboptimal, thereby increasing the likelihood of developing such aforementioned complications. Nutritional status and progress are often not adequately documented in the medical record. It can be difficult to tell when (or if) patients are consuming food and supplements. In addition, nutritional procedures and EHR-driven care recommendations are often lacking in the hospital.
- Similarly, nutritional care plans and patient issues are poorly communicated to post-acute facilities and PCPs. The current evidence supports the early and rapid identification of malnutrition in order to allow for timely treatment of malnutrition in the hospital. Part of the recommended process for implementing nutrition care is appropriate recognition, diagnosis, and documentation of the nutrition status of a patient in order to address their condition with an appropriate plan of care and communicate patient needs to other care providers.
- Identifying and addressing malnutrition early in the episode of care is associated with reduced lengths of stay, 30-day readmission rates, hospital-acquired conditions, and overall healthcare costs.

Clinical Guidance Support:

- The American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.) recommends the following: Nutrition support intervention is recommended for patients identified by screening and assessment as at risk for malnutrition or malnourished. (Grade Evidence C)


Suggested MQii Quality Indicators Can Also Be Used to Assess Quality Improvement across the Care Continuum

Malnutrition Care Workflow



The following slides provide full description of the proposed quality indicators

MQii Suggested Quality Indicators Align with the Malnutrition Care Workflow (1 of 3)



Malnutrition Screening


- Percentage of patients age 65 years and older admitted to hospital who received a malnutrition screening with a validated screening tool
- Percentage of patients age 65 years and older admitted to hospital who received a malnutrition screening
- Percentage of patients age 65 years and older identified as “at risk” through a malnutrition screening who had a malnutrition-risk diet order implemented within 24 hours of the completed screening
- Length of time between hospital admission and completion of malnutrition screening
- Length of time between identification of a patient age 65 years and older as “at risk” based on a malnutrition screening and implementation of a malnutrition-risk diet order, but before a nutrition assessment with a standardized tool
- Length of time between admission and implementation of a malnutrition-risk diet order in patients age 65 years and older identified as “at risk” based on a malnutrition screening, but before a nutrition assessment with a standardized tool

Nutrition Assessment

- Percentage of patients age 65 years and older identified as “at risk” for malnutrition based on a malnutrition screening who also had a completed nutrition assessment with a standardized tool
- Length of time between patients age 65 years and older identified as “at risk” for malnutrition based on a malnutrition screening and completion of a nutrition assessment using a standardized tool
- Length of time between admission and completion of a nutrition assessment with a standardized tool for patients age 65 years and older identified as “at risk” for malnutrition based on a malnutrition screening

MQii Suggested Quality Indicators Align with the Malnutrition Care Workflow (2 of 3)

Malnutrition Care Workflow



Malnutrition Diagnosis


- Percentage of patients age 65 years and older identified as malnourished with a nutrition assessment using a standardized tool who have a documented dietitian-based malnutrition diagnosis
- Percentage of patients age 65 years and older who have a documented provider medical diagnosis of malnutrition
- Percentage of patients age 65 years and older identified as malnourished with a nutrition assessment using a standardized tool who have a documented dietitian-based nutrition diagnosis and a provider medical diagnosis of malnutrition

Malnutrition Care Plan Development

- Percentage of patients age 65 years and older with a completed nutrition assessment and a documented malnutrition diagnosis who have a documented malnutrition care plan

MQii Suggested Quality Indicators Align with the Malnutrition Care Workflow (3 of 3)

Malnutrition Care Workflow



Intervention Implementation

- Percentage of patients age 65 years and older with a documented malnutrition diagnosis who had a nutrition intervention implemented
- Length of time between documented malnutrition diagnosis and implementation of a nutrition intervention for patients age 65 years and older diagnosed as malnourished
- Length of time between admission and implementation of a nutrition intervention for patients age 65 years and older diagnosed as malnourished

Monitoring/Evaluation & Discharge Planning

- Percentage of patients age 65 years and older with a malnutrition diagnosis as a result of a nutrition assessment with a standardized tool who have a malnutrition care plan included as part of their post-discharge care plan